

Catalysis of proton decay in superstring theory

Yuri M. Malyuta

*Institute for Nuclear Research, National Academy of Sciences of Ukraine
252022 Kiev, Ukraine*

E-mail: aks@d310.icyb.kiev.ua

Abstract

Theory of superstrings on Calabi-Yau manifolds is applied to investigate catalysis of proton decay.

1 Introduction

Rubacov [1] and Callan [2] have shown that grand unification theories which do not conserve baryon number predict the existence of monopole-induced proton decay. Recently it has been proved that Calabi-Yau models describe solitonic objects of superstring theory [3]. The purpose of this work is to study induced proton decay in the context of Calabi-Yau models.

During the last few years there has been substantial progress in superstring theory compactified on Calabi-Yau manifolds [4, 5]. In particular the idea of heterotic/type II string duality has proved to be crucial. New methods enable to investigate superstring vacua and phase transitions between them. This approach is very important because it gives new predictions for future experimental searches.

2 Catalysis reaction

Let us consider the instanton numbers (topological invariants of rational curves [4]) for the Calabi-Yau manifolds $X_{24}(1, 1, 2, 8, 12)_{-480}^{3,243}$ and $X_{20}(1, 1, 2, 6, 10)_{-372}^{4,190}$ recorded in Tables 1 and 2.

Table 1

(0, 0, 1)	−1	(0, 1, 1)	−1	(0, 1, 2)	−2	(0, 1, 3)	−3
(0, 1, 4)	−4	(0, 1, 5)	−5	(0, 2, 3)	−3	(0, 2, 4)	−16
(1, 0, 0)	240	(1, 0, 1)	240	(1, 1, 1)	240	(1, 1, 2)	720
(1, 1, 3)	1200	(1, 1, 4)	1680	(1, 2, 3)	1200	(2, 0, 0)	240
(2, 0, 2)	240	(2, 2, 2)	240	(3, 0, 0)	240	(3, 0, 3)	240
(4, 0, 0)	240	(5, 0, 0)	240	(6, 0, 0)	240	(0, 1, 0)	0

Table 2

(0, 0, 0, 1)	28	(0, 0, 0, 2)	−1	(0, 0, 0, 3)	0	(0, 0, 1, 0)	−1
(0, 1, 0, 0)	0	(0, 1, 1, 0)	−1	(0, 1, 2, 0)	−2	(0, 1, 3, 0)	−3
(0, 1, 4, 0)	−4	(0, 1, 5, 0)	−5	(0, 2, 3, 0)	−3	(0, 2, 4, 0)	−16
(0, 2, 5, 0)	−55	(0, 2, 6, 0)	−144	(0, 3, 4, 0)	−4	(0, 3, 4, 0)	−4
(0, 3, 5, 0)	−55	(1, 0, 0, 0)	−1	(1, 0, 0, 1)	28	(1, 0, 0, 2)	186
(1, 0, 0, 3)	28	(1, 0, 0, 4)	−1				

The calculation of these numbers was done using the computer program INSTANTON. From Tables 1 and 2 we infer the following relation between the instanton numbers

$$n_{a,b,c} = \sum_k n_{a,b,c,k}. \quad (1)$$

Relation (1) describes the phase transition between solitonic objects with the spectra presented in Table 3.

Table 3

Calabi-Yau	spectrum	gauge group	rank of group
$X_{24}(1, 1, 2, 8, 12)_{-480}^{3,243}$	244 1	$U(1)^4$	4
$X_{20}(1, 1, 2, 6, 10)_{-372}^{4,190}$	28 2 + 191 1	$SU(2) \times U(1)^4$	5

These spectra are calculated by using the technique of instanton numbers [5]:

$$\frac{240}{4+1} = -1 + \frac{\boxed{28}}{\boxed{244}+1} + 186 + \frac{\boxed{28}}{\boxed{191}} - 1 \quad (2)$$

We conclude from (2) that the solitonic object $Sol = (244 \text{ } \mathbf{1})$ consists of 244 singlet hypermultiplets, while the solitonic object $Sol^* = (28 \text{ } \mathbf{2} + 191 \text{ } \mathbf{1})$ consists of 28 doublet hypermultiplets and 191 singlet hypermultiplets; moreover, there exists the phase transition

$$Sol^* \rightarrow Sol + H, \quad (3)$$

where H is the singlet hypermultiplet.

Combining (3) with the quark diagram of proton decay we construct the diagram of the catalysis

$$\begin{array}{ccc}
 p \equiv \equiv \equiv \bigcirc & \longrightarrow & e^+ \\
 & \Downarrow H & \\
 Sol^* \implies \bigcirc & \implies & Sol
 \end{array} \tag{4}$$

3 Estimates of bounds

Reaction (4) has the energy emission $\varepsilon \simeq m_H$ where we equate m_H with the Higgs mass 1000 GeV [6]. To estimate the cross section of reaction (4) let us apply the Okun trick [7]. This trick is demonstrated in Table 4 where $\alpha_{12} = g_1 g_2$, g_1 and g_2 are effective coupling constants which represent the vertices of the diagram (4).

Table 4

Amplitude of reaction (4)	Probability of reaction (4)	Cross section of reaction (4)
$A = \alpha_{12} m_H^{-2}$	$\omega = A^2 \varepsilon^5 = \alpha_{12}^2 m_H$	$\sigma = \omega^{-2} = (\alpha_{12}^2 m_H)^{-2}$

Identifying in the formula for σ the constant α_{12} with the grand unification constant $\alpha_{GU} \simeq \frac{1}{40}$ [7] we obtain the upper bound of the catalysis cross section $\sigma \simeq 10^{-28} cm^2$.

References

- [1] V.A. Rubacov, JETP Lett. 33 (1981) 644.
- [2] C.G. Callan, Phys.Rev. D26 (1982) 2058.
- [3] G. Aldazabal, A. Font, L.E. Ibáñez and F. Quevedo, hep-th/9510093.
- [4] P. Berglund, S. Katz and A. Klemm, Nucl.Phys. B456 (1995) 153.
- [5] P. Berglund, S. Katz, A. Klemm and P. Mayr, hep-th/9605154.
- [6] Review of Particle Physics, Phys.Rev. D54, N 1, Part I, 1996.
- [7] L.B. Okun, Leptons and quarks (Nauka, Moscow 1981).